

VITAMIN C

Feed additives for animals in powder

COMPOSITION:

Vitamins:

- Vitamin C

Nutritional supplements:

- Sucrose
- Corn starch

USE:

Vitamin C is the strongest antioxidant among vitamins and has many roles in the organism: reductant agent in living organisms, important role in feed absorption (calcium, iron, folic acid), production of collagen, production and synthesis of steroid hormones – anti-inflammatory effect and treatment of functional sterility, synthesis of neurotransmitters, involved in carbohydrate metabolism, improves the physiological state of blood vessels, supports normal capillary function, increases the resistance towards infections, speeds up the blood coagulation and is essential for wound healing. Stress, greater physical labor, contagious diseases, gestation and lactation increase the need of this vitamin.

Vitamin C successfully helps in different diseases, hemorrhagic diathesis and during convalescence.

TARGET SPECIES: All animal species

DOSAGE:

It is administered orally after dissolving in drinking water or stirring in food for 3-5 days.

Pigs: 5g / 10l drinking water or through food 1kg per 1 ton of food

Fattening chickens, turkeys: 5-10 g / 10 liters of drinking water

Carrier hens: 10-15 g / 10 l drinking water

Pigeons and rabbits: 5-10 g / 10 l water

Horses and cattle: 5- 10 g / s

CONTRAINDICATIONS: None described

ADVERSE REACTION: None

WITHDRAWAL PERIOD: None

SPECIAL PRECAUTIONS: Store protected from humidity and light, on a temperature up to 25°C. Keep away from children. Keep in closed containers.

WAY OF ISSUE: Without a prescription

PRESENTATIONS: PE bottles, net 1 L

ANIMAL CONTROL FACILITY NUMBER: α RS-31029

PRODUCER NAME AND HEADQUARTERS: Vetplanet doo, Vrsac, Vojvodjanskih brigada 16, Serbia